

2D Isotropic Oscillator

$$H = \frac{p_x^2 + p_y^2}{2\mu} + \frac{1}{2} \mu \omega^2 (x^2 + y^2)$$

$\mu \rightarrow$ mass
 $\omega \rightarrow$ frequency

This Hamiltonian is rotationally invariant. Check if $[H, L_z] = 0$.

Due to $[H, L_z] = 0$, \hat{H} and \hat{L}_z share common eigenstate.

In cylindrical coordinates, eigenstate can be assumed as ----- (1)

$$\psi_E(\rho, \varphi) = R(\rho) \Phi(\varphi)$$

Note that $L_z |\Phi(\varphi)\rangle = m\hbar |\Phi(\varphi)\rangle$, where $\Phi_m(\varphi) = \frac{e^{im\varphi}}{\sqrt{2\pi}}$.

- $V(x, y) = \frac{1}{2} \mu \omega^2 (x^2 + y^2)$

In cylindrical-polar coordinates

$$V(\rho, \varphi) = \frac{1}{2} m \omega^2 \rho^2 = V(\rho)$$

- Schrodinger equation

$$\left[-\frac{\hbar^2}{2\mu} \left(\frac{\partial^2}{\partial \rho^2} + \frac{1}{\rho} \frac{\partial}{\partial \rho} + \frac{1}{\rho^2} \frac{\partial^2}{\partial \varphi^2} \right) + V(\rho) \right] \psi_E(\rho, \varphi) = E \psi_E(\rho, \varphi)$$

Substitute from Eq (1), and use

$$\frac{\partial^2}{\partial \varphi^2} \Phi_m(\varphi) = -m^2 \Phi_m(\varphi)$$

Then, we get the RADIAL EQUATION:

$$\left[-\frac{\hbar^2}{2\mu} \left(\frac{d^2}{d\rho^2} + \frac{1}{\rho} \frac{d}{d\rho} - \frac{m^2}{\rho^2} \right) + V(\rho) \right] R(\rho) = E R(\rho) \text{ --- (2)}$$

We already know the solution to angular part, i.e, $\Phi_m(\varphi)$.

We have to solve Eq (2) to find the radial eigenfunction $R(\rho)$.

- First determine the limiting cases:

As $\rho \rightarrow 0$:

Eq (2) can be rewritten as :

$$\left[\frac{d^2}{dp^2} + \frac{1}{p} \frac{d}{dp} - \frac{m^2}{p^2} - \frac{\mu^2 \omega^2}{2\hbar^2} p^2 + \frac{2\mu}{\hbar^2} E \right] R(p) = 0$$

As $p \rightarrow 0$, $p^2 \rightarrow 0$. But $\frac{1}{p^2} \gg 1$. So $\frac{2\mu}{\hbar^2} E$ can be neglected.

$$\therefore \text{As } p \rightarrow 0 : \left(\frac{d^2}{dp^2} + \frac{1}{p} \frac{d}{dp} - \frac{m^2}{p^2} \right) R(p) = 0$$

We can verify that $R(p) = p^{\pm m}$ is a valid solution.

However p^{-m} diverges. So, we have $R(p) \xrightarrow{p \rightarrow 0} p^{|m|}$.

$$\text{As } p \rightarrow \infty : \left(\frac{d^2}{dp^2} - \frac{\mu^2 \omega^2}{\hbar^2} p^2 \right) R(p) = 0$$

$$\text{Solution is } R(p) \xrightarrow{p \rightarrow \infty} e^{-\frac{\mu \omega}{2\hbar} p^2} p^\beta \quad (\beta \rightarrow \text{constant})$$

$$\therefore \text{We assume that } R(p) = p^{|m|} e^{-\frac{\mu \omega}{2\hbar} p^2} U(p)$$

• Now, make p dimensionless with the change of variables

$$y = \sqrt{\frac{\mu \omega}{\hbar}} p \quad \text{and} \quad \epsilon = \frac{E}{\hbar \omega}$$

Radial equation (2) for $U(y)$ becomes

$$\frac{d^2 U}{dy^2} + \left[\left(\frac{2|m|+1}{y} \right) - 2y \right] \frac{dU}{dy} + (2\epsilon - 2|m| - 2)U = 0$$

• Use power series method to solve this eqn:

$$U(y) = \sum_{r=0}^{\infty} c_r y^r$$

Obtain the recurrence relation:

$$C_{r+2} = \frac{2r - (2\epsilon - 2|m| - 2)}{(r+2)[2|m| + r + 2]} C_r$$

This series must terminate at finite r if $y \rightarrow \infty$ behaviour should come out right.

$$2\epsilon - 2|m| - 2 = 2r$$

$$\Rightarrow \epsilon = r + |m| + 1$$

NOTE: r must be an even number, i.e., $r = 2k$
 $\therefore \frac{E}{\hbar\omega} = (2k + |m| + 1) \Rightarrow E_k = (2k + |m| + 1) \hbar\omega$
 $k = 0, 1, 2, \dots$
 \downarrow radial quantum number


- put $n = 2k + |m|$. $E_n = (n+1) \hbar\omega$
- For a given n , what are allowed values of $|m|$
- " " " , degeneracy is $(n+1)$.

k	m	n	E_n
0	0	0	$1\hbar\omega$
0	1	1	$2\hbar\omega$
0	-1	1	$2\hbar\omega$
0	2	2	$3\hbar\omega$
0	-2	2	$3\hbar\omega$
1	0	2	$3\hbar\omega$

Exercise: Show in general that for a given value of n , degree of degeneracy is $n+1$.

Eigenfunctions:

$$\psi_0(r, \varphi) = \frac{C_0}{\sqrt{2\pi}} e^{-\frac{\mu\omega}{2\hbar} r^2}$$




normalise to get C_0 :

$$C_0 = \sqrt{2\mu\omega/\hbar}$$

Agrees with cartesian coordinate result.

1] has $k=0$ and $m=\pm 1$

$$\psi_{n=1, m=1}(r, \varphi) = \frac{C_0}{\sqrt{2\pi}} \sqrt{\frac{\mu\omega}{\hbar}} r e^{-\frac{\mu\omega}{2\hbar} r^2} e^{i\varphi}$$



$$\psi_{n=1, m=-1}(r, \varphi) = \dots e^{-i\varphi}$$

$$C_0 = \sqrt{2\mu\omega/\hbar}$$

cartesian coordinates:

$$\psi_{10} = \frac{\sqrt{2\mu\omega}}{\hbar\sqrt{\pi}} e^{-\frac{\mu\omega}{2\hbar} r^2} r \cos \varphi$$

$$\psi_{01} = \dots e^{-\frac{\mu\omega}{2\hbar} r^2} r \sin \varphi$$

use $r^2 = x^2 + y^2$
 $x = r \cos \varphi$
 $y = r \sin \varphi$

$$\psi_{n=1, m=1} = \frac{1}{\sqrt{2}} (\psi_{10} + i\psi_{01})$$

$$\psi_{n=1, m=-1} = \frac{1}{\sqrt{2}} (\psi_{10} - i\psi_{01})$$