

PHY 4154

Nuclear and Particle Physics

Sourabh Dube

August 2025

Nucleon Isospin

- ▶ Heisenberg noticed the similarities in masses between neutron and proton. ($m_p = 938.28 \text{ MeV}/c^2$, $m_n = 939.57 \text{ MeV}/c^2$).
- ▶ “Their charge is different, but aside from this they are identical”
- ▶ Charge independence of hadronic force.
- ▶ What gives rise to this symmetry?

Nucleon Isospin

We begin by noticing that when $s = \frac{1}{2}$, then $|\frac{1}{2}, \frac{1}{2}\rangle$ and $|\frac{1}{2}, -\frac{1}{2}\rangle$ are two states of the same particle.

Nucleon Isospin

We begin by noticing that when $s = \frac{1}{2}$, then $|\frac{1}{2}, \frac{1}{2}\rangle$ and $|\frac{1}{2}, -\frac{1}{2}\rangle$ are two states of the same particle.

Let there be a particle called nucleon with a property called Isospin, I . It has 3 components, I_1, I_2, I_3 .

Nucleon Isospin

We begin by noticing that when $s = \frac{1}{2}$, then $|\frac{1}{2}, \frac{1}{2}\rangle$ and $|\frac{1}{2}, -\frac{1}{2}\rangle$ are two states of the same particle.

Let there be a particle called nucleon with a property called Isospin, I . It has 3 components, I_1, I_2, I_3 .

In analogy to spin, nucleon has $I = \frac{1}{2}$, and we have two possible states

$$|\frac{1}{2}, \frac{1}{2}\rangle = \text{proton} ; \quad |\frac{1}{2}, -\frac{1}{2}\rangle = \text{neutron}$$

$$\begin{pmatrix} 1 \\ 0 \end{pmatrix} = \text{proton} ; \quad \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \text{neutron}$$

Nucleon Isospin

We begin by noticing that when $s = \frac{1}{2}$, then $|\frac{1}{2}, \frac{1}{2}\rangle$ and $|\frac{1}{2}, -\frac{1}{2}\rangle$ are two states of the same particle.

Let there be a particle called nucleon with a property called Isospin, I . It has 3 components, I_1, I_2, I_3 .

In analogy to spin, nucleon has $I = \frac{1}{2}$, and we have two possible states

$$|\frac{1}{2}, \frac{1}{2}\rangle = \text{proton} ; \quad |\frac{1}{2}, -\frac{1}{2}\rangle = \text{neutron}$$

$$\begin{pmatrix} 1 \\ 0 \end{pmatrix} = \text{proton} ; \quad \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \text{neutron}$$

Note that components are not in regular space, but in abstract 'isospin' space.

Nucleon Isospin

The generators of isospin are $I_i = \frac{1}{2}\tau_i$

$$\tau_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \tau_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \tau_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

Recall the Pauli spin matrices, with $\hat{\mathbf{S}} = \left(\frac{\hbar}{2}\right) \boldsymbol{\sigma}$

$$\sigma_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

Nucleon Isospin

Charge independence of nuclear force $\Leftrightarrow$ Strong interactions are invariant under rotations in isospin space.

Isospin is conserved in all strong interactions.

Nucleon Isospin

Charge independence of nuclear force $\Leftrightarrow$ Strong interactions are invariant under rotations in isospin space.

Isospin is conserved in all strong interactions.

Thus if $H = H_{em} + H_{had}$, then we have $[H_{had}, I] = 0$.

But $[H_{had} + H_{em}, I] \neq 0$.

Nucleon Isospin

Charge independence of nuclear force $\Leftrightarrow$ Strong interactions are invariant under rotations in isospin space.

Isospin is conserved in all strong interactions.

Thus if $H = H_{em} + H_{had}$, then we have $[H_{had}, I] = 0$.

But $[H_{had} + H_{em}, I] \neq 0$.

However we know $[H_{had} + H_{em}, Q] = 0$.

We could define charge for $|I, I_3\rangle$ as $q = e(I_3 + \frac{1}{2})$.

This gives $q_p = 1$ ($I_3 = \frac{1}{2}$), and $q_n = 0$ ($I_3 = -\frac{1}{2}$).

We have $[H_{had} + H_{em}, I_3] = 0$.

Pion Isospin

We have $[H_{had}, I] = 0$. All particles with same I have same hadronic properties. Different members of an isospin multiplet are in essence same particle with different orientations in isospin space.

Pion Isospin

We have $[H_{had}, I] = 0$. All particles with same I have same hadronic properties. Different members of an isospin multiplet are in essence same particle with different orientations in isospin space.

Consider the pion-exchange theory of Yukawa to explain strong forces
 $N \rightarrow N' + \pi$.

Isospin is conserved, and nucleon isospin $I = \frac{1}{2}$. What possible values can pion isospin have?

Pion Isospin

We have $[H_{had}, I] = 0$. All particles with same I have same hadronic properties. Different members of an isospin multiplet are in essence same particle with different orientations in isospin space.

Consider the pion-exchange theory of Yukawa to explain strong forces $N \rightarrow N' + \pi$.

Isospin is conserved, and nucleon isospin $I = \frac{1}{2}$. What possible values can pion isospin have?

$$\frac{1}{2} \rightarrow \frac{1}{2} \oplus I_\pi$$

It can be 0 or 1 .

If $I = 0$, there would be one pion. But there are three pions ($\pi^\pm, \pi^0$). Thus for pions we assign $I = 1$.


$$\pi^+ = |1, 1\rangle, \quad \pi^0 = |1, 0\rangle, \quad \pi^- = |1, -1\rangle$$

Δ Isospin

For Δ 's, $I = \frac{3}{2}$

$$\Delta^{++} = \left| \frac{3}{2}, \frac{3}{2} \right\rangle, \quad \Delta^+ = \left| \frac{3}{2}, \frac{1}{2} \right\rangle$$

$$\Delta^0 = \left| \frac{3}{2}, -\frac{1}{2} \right\rangle, \quad \Delta^- = \left| \frac{3}{2}, -\frac{3}{2} \right\rangle$$




Δ Isospin

For Δ 's, $I = \frac{3}{2}$

$$\Delta^{++} = \left| \frac{3}{2}, \frac{3}{2} \right\rangle, \quad \Delta^+ = \left| \frac{3}{2}, \frac{1}{2} \right\rangle$$

$$\Delta^0 = \left| \frac{3}{2}, -\frac{1}{2} \right\rangle, \quad \Delta^- = \left| \frac{3}{2}, -\frac{3}{2} \right\rangle$$




Essentially, to assign I , take number of particles in a multiplet. They have I_3 from $-I$ to I , and so number $= 2I + 1$.

Δ Isospin

For Δ 's, $I = \frac{3}{2}$

$$\Delta^{++} = \left| \frac{3}{2}, \frac{3}{2} \right\rangle, \quad \Delta^+ = \left| \frac{3}{2}, \frac{1}{2} \right\rangle$$


$$\Delta^0 = \left| \frac{3}{2}, -\frac{1}{2} \right\rangle, \quad \Delta^- = \left| \frac{3}{2}, -\frac{3}{2} \right\rangle$$



Essentially, to assign I , take number of particles in a multiplet. They have I_3 from $-I$ to I , and so number = $2I + 1$.

For hadrons composed of only the u, d, s quarks, we have the Gell-Mann–Nishijima formula $Q = I_3 + \frac{1}{2}(B + S)$. B is baryon number, and S is strangeness.


Isospin



$Q = I_3 + \frac{1}{2}(B + S)$, what is Σ and Ξ isospins?

Σ^+	uus	Σ^0	uds
Σ^-	dds		
Ξ^0	uss	Ξ^-	dss

Isospin



$Q = I_3 + \frac{1}{2}(B + S)$, what is Σ and Ξ isospins?

Σ^+	uus	Σ^0	uds
Σ^-	dds		
Ξ^0	uss	Ξ^-	dss

In context of quark model, it can follow from u, d being a isospin doublet ($I = \frac{1}{2}$), and s having isospin zero. (All other flavors also get isospin zero, as do leptons and mediators).

Implications of Isospin

Consider adding the isospins of p and n , i.e. a state with two nucleons (each with $I = \frac{1}{2}$), will result in

$$|1, 1\rangle = pp$$

$$|1, 0\rangle = \left(\frac{1}{\sqrt{2}}\right) (pn + np)$$

$$|1, -1\rangle = nn$$

$$|0, 0\rangle = \left(\frac{1}{\sqrt{2}}\right) (pn - np)$$

where by pp we mean the state $|\frac{1}{2}, \frac{1}{2}\rangle |\frac{1}{2}, \frac{1}{2}\rangle$ and so on

Implications of Isospin

Consider adding the isospins of p and n , i.e. a state with two nucleons (each with $I = \frac{1}{2}$), will result in

$$|1, 1\rangle = pp$$

$$|1, 0\rangle = \left(\frac{1}{\sqrt{2}}\right) (pn + np)$$

$$|1, -1\rangle = nn$$

$$|0, 0\rangle = \left(\frac{1}{\sqrt{2}}\right) (pn - np)$$

where by pp we mean the state $|\frac{1}{2}, \frac{1}{2}\rangle |\frac{1}{2}, \frac{1}{2}\rangle$ and so on

Experimentally we only see one bound state of p and n (and no pp or nn bound states). Thus deuteron must be isosinglet (strong attraction in $I = 0$ channel, not in $I = 1$ channel). The deuteron isospin is thus 0.

Nucleon nucleon scattering

Consider the following processes

(a) $p + p \rightarrow d + \pi^+$

(b) $p + n \rightarrow d + \pi^0$

(c) $n + n \rightarrow d + \pi^-$

Deuteron has $I = 0$.

On RHS isospin states are defined by pion isospin:

$|1, 1\rangle, |1, 0\rangle, |1, -1\rangle$.

Nucleon nucleon scattering

Consider the following processes

(a) $p + p \rightarrow d + \pi^+$

(b) $p + n \rightarrow d + \pi^0$

(c) $n + n \rightarrow d + \pi^-$

Deuteron has $I = 0$.

On RHS isospin states are defined by pion isospin:

$$|1, 1\rangle, |1, 0\rangle, |1, -1\rangle.$$

On LHS (see prev slide), the pp and nn states are $|1, 1\rangle$, and $|1, -1\rangle$.

The pn state is $\sqrt{\frac{1}{2}}(|1, 0\rangle + |0, 0\rangle)$.

Nucleon nucleon scattering

Consider the following processes

(a) $p + p \rightarrow d + \pi^+$

(b) $p + n \rightarrow d + \pi^0$

(c) $n + n \rightarrow d + \pi^-$

Deuteron has $I = 0$.

On RHS isospin states are defined by pion isospin:

$|1, 1\rangle, |1, 0\rangle, |1, -1\rangle$.

On LHS (see prev slide), the pp and nn states are $|1, 1\rangle$, and $|1, -1\rangle$.

The pn state is $\sqrt{\frac{1}{2}}(|1, 0\rangle + |0, 0\rangle)$.

Since isospin is conserved, only the $I = 1$ state will contribute. Thus the scattering amplitudes are in the ratio $\mathcal{M}_a : \mathcal{M}_b : \mathcal{M}_c = 1 : \sqrt{\frac{1}{2}} : 1$.

The cross section, σ , goes like $|\mathcal{M}|^2$, and thus are in the ratio $\sigma_a : \sigma_b : \sigma_c = 2 : 1 : 2$.

Cross section σ is the rate at which a process occurs, and is measured in units of area (more on this later).

Pion-nucleon scattering

Pion-nucleon scattering processes ($\pi N \rightarrow \pi N$).

Elastic processes

(a) $\pi^+ + p \rightarrow \pi^+ + p$

(c) $\pi^- + p \rightarrow \pi^- + p$

(e) $\pi^0 + n \rightarrow \pi^0 + n$

(b) $\pi^0 + p \rightarrow \pi^0 + p$

(d) $\pi^+ + n \rightarrow \pi^+ + n$

(f) $\pi^- + n \rightarrow \pi^- + n$

Charge-exchange processes

(g) $\pi^+ + n \rightarrow \pi^0 + p$

(i) $\pi^0 + n \rightarrow \pi^- + p$

(h) $\pi^0 + p \rightarrow \pi^+ + n$

(j) $\pi^- + p \rightarrow \pi^0 + n$

Pion-nucleon scattering

Pion-nucleon scattering processes ($\pi N \rightarrow \pi N$).

Elastic processes

$$(a) \pi^+ + p \rightarrow \pi^+ + p$$

$$(c) \pi^- + p \rightarrow \pi^- + p$$

$$(e) \pi^0 + n \rightarrow \pi^0 + n$$

$$(b) \pi^0 + p \rightarrow \pi^0 + p$$

$$(d) \pi^+ + n \rightarrow \pi^+ + n$$

$$(f) \pi^- + n \rightarrow \pi^- + n$$

Charge-exchange processes

$$(g) \pi^+ + n \rightarrow \pi^0 + p$$

$$(i) \pi^0 + n \rightarrow \pi^- + p$$

$$(h) \pi^0 + p \rightarrow \pi^+ + n$$

$$(j) \pi^- + p \rightarrow \pi^0 + n$$

The pion has $I = 1$, and the nucleon has $I = \frac{1}{2}$

thus the total isospin can be $3/2$ or $1/2$.

There are only two distinct scattering amplitudes here

(strong processes invariant under rotations in isospin space),

lets call these as $\mathcal{M}_3$ for $I = \frac{3}{2}$ and $\mathcal{M}_1$ for $I = \frac{1}{2}$.

Pion-nucleon scattering

$$\begin{aligned}\pi^+ + p &: |1, 1\rangle \left| \frac{1}{2}, \frac{1}{2} \right\rangle &= \left| \frac{3}{2}, \frac{3}{2} \right\rangle \\ \pi^0 + p &: |1, 0\rangle \left| \frac{1}{2}, \frac{1}{2} \right\rangle &= \sqrt{\frac{2}{3}} \left| \frac{3}{2}, \frac{1}{2} \right\rangle - \sqrt{\frac{1}{3}} \left| \frac{1}{2}, \frac{1}{2} \right\rangle \\ \pi^- + p &: |1, -1\rangle \left| \frac{1}{2}, \frac{1}{2} \right\rangle &= \sqrt{\frac{1}{3}} \left| \frac{3}{2}, -\frac{1}{2} \right\rangle - \sqrt{\frac{2}{3}} \left| \frac{1}{2}, -\frac{1}{2} \right\rangle \\ \pi^+ + n &: |1, 1\rangle \left| \frac{1}{2}, -\frac{1}{2} \right\rangle &= \sqrt{\frac{1}{3}} \left| \frac{3}{2}, \frac{1}{2} \right\rangle + \sqrt{\frac{2}{3}} \left| \frac{1}{2}, \frac{1}{2} \right\rangle \\ \pi^0 + n &: |1, 0\rangle \left| \frac{1}{2}, -\frac{1}{2} \right\rangle &= \sqrt{\frac{2}{3}} \left| \frac{3}{2}, -\frac{1}{2} \right\rangle + \sqrt{\frac{1}{3}} \left| \frac{1}{2}, -\frac{1}{2} \right\rangle \\ \pi^- + n &: |1, -1\rangle \left| \frac{1}{2}, -\frac{1}{2} \right\rangle &= \left| \frac{3}{2}, -\frac{3}{2} \right\rangle\end{aligned}$$

Pion-nucleon scattering

Pion-nucleon scattering processes ($\pi N \rightarrow \pi N$).

$$\begin{aligned}
 \text{(a)} \quad \pi^+ + p &\rightarrow \pi^+ + p & \left| \frac{3}{2}, \frac{3}{2} \right\rangle &\rightarrow \left| \frac{3}{2}, \frac{3}{2} \right\rangle \\
 \text{(f)} \quad \pi^- + n &\rightarrow \pi^- + n & \left| \frac{3}{2}, -\frac{3}{2} \right\rangle &\rightarrow \left| \frac{3}{2}, -\frac{3}{2} \right\rangle \\
 \text{(c)} \quad \pi^- + p &\rightarrow \pi^- + p & \sqrt{\frac{1}{3}} \left| \frac{3}{2}, -\frac{1}{2} \right\rangle - \sqrt{\frac{2}{3}} \left| \frac{1}{2}, -\frac{1}{2} \right\rangle &\rightarrow \sqrt{\frac{1}{3}} \left| \frac{3}{2}, -\frac{1}{2} \right\rangle - \sqrt{\frac{2}{3}} \left| \frac{1}{2}, -\frac{1}{2} \right\rangle \\
 \text{(b)} \quad \pi^0 + p &\rightarrow \pi^0 + p & \sqrt{\frac{2}{3}} \left| \frac{3}{2}, \frac{1}{2} \right\rangle - \sqrt{\frac{1}{3}} \left| \frac{1}{2}, \frac{1}{2} \right\rangle &\rightarrow \sqrt{\frac{2}{3}} \left| \frac{3}{2}, \frac{1}{2} \right\rangle - \sqrt{\frac{1}{3}} \left| \frac{1}{2}, \frac{1}{2} \right\rangle \\
 \text{(j)} \quad \pi^- + p &\rightarrow \pi^0 + n & \sqrt{\frac{1}{3}} \left| \frac{3}{2}, -\frac{1}{2} \right\rangle - \sqrt{\frac{2}{3}} \left| \frac{1}{2}, -\frac{1}{2} \right\rangle &\rightarrow \sqrt{\frac{2}{3}} \left| \frac{3}{2}, -\frac{1}{2} \right\rangle + \sqrt{\frac{1}{3}} \left| \frac{1}{2}, -\frac{1}{2} \right\rangle
 \end{aligned}$$

We have $\mathcal{M}_a = \mathcal{M}_f = \mathcal{M}_3$

$$\mathcal{M}_c = \frac{1}{3}\mathcal{M}_3 + \frac{2}{3}\mathcal{M}_1$$

$$\mathcal{M}_b = \frac{2}{3}\mathcal{M}_3 + \frac{1}{3}\mathcal{M}_1$$

$$\mathcal{M}_j = \frac{\sqrt{2}}{3}\mathcal{M}_3 - \frac{\sqrt{2}}{3}\mathcal{M}_1$$

$$\sigma_a : \sigma_c : \sigma_j = 9|\mathcal{M}_3|^2 : |\mathcal{M}_3 + 2\mathcal{M}_1|^2 : 2|\mathcal{M}_3 - \mathcal{M}_1|^2$$

Quick data analysis concept

Consider doing this scattering experiment (with $\pi^\pm$ and protons).

Let me explain one quick concept first.

Suppose you have a decay of particle $A \rightarrow B + C$

Evidently conservation of energy-momentum $\Rightarrow p_A = p_B + p_C$
(where p are 4-vectors!)

Experimentally we measure p_B and p_C , and thus calculate $(p_B + p_C)^2$,
but this is just p_A^2 and $p_A^2 = m_A^2$
i.e. $\sqrt{(p_B + p_C)^2} = m_A$

Thus by measuring the 4-vectors of the daughters (B, C), we are able to infer the mass of the mother (A). The LHS calculated quantity is called the **invariant mass of B and C** and is calculated from the observed/measured 4-vectors of B and C .

Invariant mass

Given two 4-vectors p and q , with components (E_p, p_x, p_y, p_z) and (E_q, q_x, q_y, q_z) respectively,


the invariant mass is defined by first adding the two vectors

$$r = p + q = (E_p + E_q, p_x + q_x, p_y + q_y, p_z + q_z)$$

and then calculating

$$r^2 = [(E_p + E_q)^2 - (p_x + q_x)^2 - (p_y + q_y)^2 - (p_z + q_z)^2]$$

Invariant mass example



The invariant mass spectrum of two charged leptons.
i.e either two muons ($\mu^+\mu^-$) or two electrons (e^+e^-)

Pion-nucleon scattering

Consider doing this scattering experiment (with $\pi^\pm$ and protons).

Given that measuring total cross section is easier, we can measure the total cross section of $(\pi^+ + p)$ vs $(\pi^- + p)$

(a) $\pi^+ + p \rightarrow \pi^+ + p$

(c) $\pi^- + p \rightarrow \pi^- + p$

(j) $\pi^- + p \rightarrow \pi^0 + n$

Pion-nucleon scattering

Consider doing this scattering experiment (with $\pi^\pm$ and protons).

Given that measuring total cross section is easier, we can measure the total cross section of $(\pi^+ + p)$ vs $(\pi^- + p)$

$$(a) \pi^+ + p \rightarrow \pi^+ + p$$

$$(c) \pi^- + p \rightarrow \pi^- + p$$

$$(j) \pi^- + p \rightarrow \pi^0 + n$$

$$\sigma_a : \sigma_c : \sigma_j =$$

$$9|\mathcal{M}_3|^2 : |\mathcal{M}_3 + 2\mathcal{M}_1|^2 : 2|\mathcal{M}_3 - \mathcal{M}_1|^2$$

At a CM energy of 1232 MeV, we get the

Δ 'resonance' ($I = \frac{3}{2}$), thus $\mathcal{M}_3 \gg \mathcal{M}_1$ and




$$\sigma_a : \sigma_c : \sigma_j = 9 : 1 : 2$$

$$\frac{\sigma_{tot}(\pi^+ + p)}{\sigma_{tot}(\pi^- + p)} = 3$$

Pion-nucleon scattering

Consider doing this scattering experiment (with $\pi^\pm$ and protons).

Given that measuring total cross section is easier, we can measure the total cross section of $(\pi^+ + p)$ vs $(\pi^- + p)$




$$\sigma_a : \sigma_c : \sigma_j =$$

$$9|\mathcal{M}_3|^2 : |\mathcal{M}_3 + 2\mathcal{M}_1|^2 : 2|\mathcal{M}_3 - \mathcal{M}_1|^2$$

At a CM energy of 1232 MeV, we get the Δ 'resonance' ($I = \frac{3}{2}$), thus $\mathcal{M}_3 \gg \mathcal{M}_1$ and

$$\sigma_a : \sigma_c : \sigma_j = 9 : 1 : 2$$

$$\frac{\sigma_{tot}(\pi^+ + p)}{\sigma_{tot}(\pi^- + p)} = 3$$



Spectrum from Griffiths, quark flow diagram from A.J.Barr